ALLOWED AS OF IN 106

. 2.

Applicants: Virginia M. Litwin et al.

Serial No.: 09/891,062 Filed: June 25, 2001

Page 3

## Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

## Listing of Claims

1-39. (Canceled)

- L 40. (Currently Amended) A method of inhibiting HIV-1 infection of a CD4+ cell which comprises contacting the CD4+ cell with an amount of a monoclonal antibody or portion thereof effective to (a) specifically inhibit 67% or greater of fusion of a CD4+ PM-1 cell to a HeLa cell expressing envelope glycoprotein from HIV-1<sub>JR-FL</sub>, and (b) inhibit 18% or less of fusion of a CD4+ SUP-T1 cell to a HeLa cell expressing envelope protein from HIV-1LAI, wherein the antibody (i) does not crossreact with HIV-1 envelope glycoprotein or CD4, (ii) reacts with an antigen on the surface of a PM-1 cell having an approximate molecular weight of 44 kD, (iii) does not react with an antigen on the surface of a SUP-T1 cell, and (iv) is at least as active as monoclonal antibody PA-7 produced by the hybridoma designated PA-7 (ATCC Accession No. PTA-6638) in inhibiting fusion as recited in (a) above and less active than monoclonal antibody PA-6 produced by the hybridoma designated PA-6 (ATCC Accession No. PTA-6637) in inhibiting fusion as recited in (b) above, so as to thereby inhibit HIV-1 infection of the CD4+ cell.
- 24. (Previously Presented) The method of claim 40, wherein the monoclonal antibody is a chimeric monoclonal antibody.
- 3 4. (Previously Presented) The method of claim 40, wherein the

Applicants: Vi

Virginia M. Litwin et al.

Serial No.: Filed:

09/891,062 June 25, 2001

Page 4

monoclonal antibody is humanized.

43. (Previously Presented) The method of claim 40, wherein the monoclonal antibody is a human monoclonal antibody.

- (Previously Presented) The method of claim 40, wherein the portion of the monoclonal antibody is a single chain antibody or an antigen binding fragment.
- (Previously Presented) The method of claim 40, wherein the monoclonal antibody is labeled with a detectable marker.
- 7 %. (Previously Presented) The method of claim 46, wherein the detectable marker is a radioactive isotope, enzyme, dye or biotin.
- (Previously Presented) The method of claim 40, wherein the CD4+ cell is present in the subject and the contacting is effected by administering the monoclonal antibody or portion thereof to the subject.